



**Connecticut**  
**Department of Energy &  
Environmental Protection**

**Field Visit to the Scodon Drive Property in Ridgefield, CT  
Post Visit Report for 34 Acres**

**Present Parties:** Conservation Commission: Tony Markert and David Beers (Western District Service Forester) on 1/7/2026 from 12:30-3

**Stewardship Objectives**

1. Maintain properties in an ecologically sound manner
2. Maintain properties for the recreational benefit of the town



---

## PROPERTY OVERVIEW

---

The property has public road frontage on George Washington Highway to the north, Pheasant Lane to the west, and Scodon Drive to the south. The only good access is at a short gated access drive to an Aquarion well on Scodon Drive. It is surrounded by primarily residential development, with a few patches of forest and farmland. Ridgefield's Old Trolley Preserve abuts the property to the north. This is a mainly developed landscape, with a few small blocks of forest and farmland.

This forest is part of a small core forest block having less than 250 acres of contiguous forest. Core forests are large tracts of unbroken forest that provide a much more stable home for plant and animal species, thereby protecting biodiversity. They are forested areas surrounded by more forested areas.

The CT DEEP Natural Diversity Database (NDDDB) does **not** record occurrences of threatened or endangered species on or near this property. Maps showing wetland and farmland soils are attached to this report. This property is in the Still River and Housatonic River watershed.

In the attached 1934 air photo, the central area surrounded by stonewalls was fields, and the remainder of the property was forest. This makes sense because the central area has prime farmland.

Please see the attached appendix for more information about your forest's history, future, and general recommendations. It also includes an appendix of Latin names for the tree and shrub species.



---

## FOREST VEGETATION (~34 ACRES)

---

### Tree Cover

#### Most Common

Red Oak  
White Oak  
Hickory  
Tulip Poplar  
Red Maple  
Sugar Maple  
Black Birch  
Yellow Birch  
Black Locust

#### Less Common

Beech  
Black Oak  
Scarlet Oak  
Black Cherry  
Sassafras  
Blackgum  
Slippery Elm  
Basswood

This forest features a diverse mix of tree species and sizes, growing in site conditions that vary with slope position, land-use history, and soil type. Over a third of the property is forested wetlands. The central hill, surrounded by stone walls, is a younger forest with more locust trees, tulip poplar trees, and barberry shrubs.

### Understory

There are a few maple, beech and birch saplings. There are also some musclewood, spicebush, high-bush blueberry and greenbrier shrubs.

### Ground Cover

In areas with thick invasives, the leaf litter is degraded and thin. Other areas appear to have thick, healthy leaf litter. There is a good amount of woody material throughout the forest floor.

### Forest Health

Beech leaf disease has infected the beech trees. This microscopic nematode, originating in Asia, has recently begun spreading throughout the state, and the long-term prognosis for infected beech trees remains unclear. The emerald ash borer has killed ash trees within the past decade.

The canopy structural diversity and the good tree and shrub species diversity make this forest more resilient to future disturbances (weather, climate, pests). The large deer herd in Connecticut makes growing certain preferred deer browse, such as oak and hickory seedlings, difficult.

Patches of exotic invasive vegetation grow in the understory, including barberry shrubs, multiflora rose shrubs, privet shrubs and bittersweet vines. Some areas are thick with barberry shrubs. There are patches of bittersweet and grape vines throughout the forest. Please refer to the appendix for additional information on invasive species and their control.

### Wildlife Habitat

The mature oak and hickory trees in your forest are an excellent asset for wildlife, especially in terms of the acorns and nuts they produce. Having diverse habitat types (younger, mature, wetlands) intimately mixed, as you have, is always good. Your forest provides adequate food, water, shelter, cover, and space for much of the wildlife in the area. Please refer to the appendix for additional information on wildlife habitats.

### Carbon and Climate Resilience

This mature forest stores a large amount of carbon while sequestering more. Any forest products you produce will help

mitigate climate change. The following recommendations aim to enhance forests' resilience and adaptability to climate change. More information about forest carbon and the forest's ecological services is in the appendix.

### **Forest Vegetation - Potential Recommendations**

The tree-hanging, exotic bittersweet and grape vines reduce the vigor of mature trees by shading and choking their upper canopy, strangling stems and branches, and possibly even pulling down tree branches. I recommend cutting these vines with a handsaw or chainsaw. This cutting can be done during the growing season when the vine's energy stores are above ground. Two cuts should be made at least 4' apart to prevent vine reconnection and to create an open window where new vines have nothing to grab onto (eliminate laddering opportunities). The cut vines will not survive in shaded areas. To be extra sure, you can treat the freshly cut rooted end with an herbicide immediately after cutting – see photo of the buckthorn blaster in the appendix.



Tall Large Tulip Poplar Tree

---

## CONCLUSION

---

### Here are some possibilities for your forest:

- Contact a private forester about doing a forest stewardship plan\*
- Properly locate and mark your property boundaries (see appendix)
  - An annual property inspection that includes property boundaries
- Cut the vines
- Potential for a loop trail from the Aquarion well gate that would include a boardwalk wetland crossing
- Enjoy your forest!

\*Please consider hiring a forester to help you implement any of the recommendations in this report (see appendix). I highly recommend considering a grant to support this work: [Urban Forestry Grant Opportunities](#). Please feel free to share this report.


It is pertinent to note that this is not a stewardship plan and is not intended as a substitute for one. It is a recap report based on a brief tour of the property, without collecting any data as needed for proper planning.



Younger Central Forest

Scodon Drive Property  
Ridgefield Conservation Commission  
Ridgefield, CT  
34 Acres

Prepared by David Beers  
CT DEEP Service Forester  
1/7/2026




- Old Fencelines
- Woods Roads
- Streams
- Wetlands
- Bounds
- 10 FT Contours



0 187.5 375 750 1,125 1,500 Feet

Scodon Drive Property  
 Ridgefield Conservation Commission  
 Ridgefield, CT  
 34 Acres

Prepared by David Beers  
 CT DEEP Service Forester  
 1/7/2026




- Old Fencelines
- Woods Roads
- Streams
- Wetlands
- Bounds
- 10 FT Contours
- Wetland Soils**
- Wetland Soils
- Floodplain Soils
- Farmland Soils**
- Prime Farmland Soils
- Statewide Important Farmland Soils
- Locally Important Farmland Soils



Scodon Drive Property  
Ridgefield Conservation Commission  
Ridgefield, CT  
34 Acres

Prepared by David Beers  
CT DEEP Service Forester  
1/7/2026  
1934 Air Photo





---

**APPENDIX**

---

**FOREST HISTORY**

Between the eighteenth-century colonial settlement and the mid-nineteenth century, most of Connecticut was cleared for agriculture, leaving only a few small patches of forest. Only 25% of Connecticut was forested in 1830. Under these conditions, a muskrat was the largest animal in the woods. Turkeys, deer, bobcats, beavers, and bears were either rare or gone. Most of the land was used for livestock pasture, with only the best soils used for hay or tilled crops. Imagine a very open agricultural landscape.

During this farming period, the stonewalls were built to keep livestock out of crops and the neighbor's property. Most of these walls were topped off with piled wood and stumps to increase their height. Stonewalls were also a depository for rocks removed from cultivated land. A stonewall with many fist-sized rocks means that one side of that wall had tilled crops, where the winter freeze of bare ground would push rocks to the surface. After barbed wire became widely available in 1875, many of these walls were supplemented with wire. Barbed wire was used to corral cows and goats, but not sheep (the barbs did not hurt them). Sheep pasture used smooth-wire rectangular page fencing.

Most Connecticut hill farms were abandoned between the mid-nineteenth and early twentieth centuries. The farmers either moved west in search of better farming soils or headed to the cities for employment. Immediately after the farm's abandonment, the forest began reclaiming the land. Much of the young forestland was then repeatedly cut down to produce charcoal, which fueled industrialization. Charcoal was used in metal blast furnaces and by blacksmiths.

Small trees were cut into 4-foot lengths and carried by hand to make a circular pile about 30 feet wide and 10 feet high for charcoal making. A ditch was dug around the circumference of the pile, and the soil from the ditch covered the pile to limit the amount of oxygen in the smoldering pile. Once the low-oxygen burn was completed in two weeks, the almost pure carbon charcoal was removed for transport to market. Charcoal produces the fire needed for metalworking.

While this charcoal-making process had been occurring since settlement, it reached its peak between 1880 and 1920. At that time, much of the landscape was cut multiple times, with patches of smoke rising from active charcoal mounds across the hills. By about 1925, cheaper coal led to the decline of charcoal production, and the forest began to grow back. The repetitive cutting of young trees for charcoal encouraged the proliferation of oak trees. Of all the tree species, oak responded most favorably to repetitive cutting. This, along with frequent wildfires, contributed to the development of the oak-dominated forest we see today.

The 1934 map is attached. Please note that you need to mentally adjust the map, as the map scale projection does not exactly match what we use today. To see what ancestral homeland existed on your property before settlement, please visit [Native-Land.ca](http://Native-Land.ca), and type in your address.

**FOREST FUTURE**

Active forest management can nudge a forest in different directions by manipulating which trees continue to grow and how much sunlight the forest floor receives by creating canopy openings of various sizes and shapes. For example, we can nudge the future forest towards oak by leaving oaks to grow and produce acorns, creating canopy openings of sufficient size to bring in the sunlight young oaks need to grow, and hunting the deer that eat young oak trees. Without these manipulations and significant natural disturbances (wind, ice, pests), the forest will gradually transition to shade-tolerant trees not eaten by deer (hemlock, beech, black birch and red maple).

## GENERAL RECOMMENDATIONS FOR FOREST STEWARDSHIP

### Forest Protection

For long-term protection, a conservation easement on part of the property could be donated to a local land trust, preventing development in perpetuity. The easement's value may be income tax-deductible over several years and provide estate tax benefits.

1. Please contact your local land trusts for more information using this link: [Find A Land Trust - Connecticut Land Conservation Council \(ctconservation.org\)](https://www.ctconservation.org/).
2. Please contact NRCS about HFRP (Healthy Forest Reserve Program) for conservation easement funding using this link: [NRCS Accepting Applications for HFRP Program | Natural Resources Conservation Service](https://www.nrcs.usda.gov/programs/healthy-forest-reserve-program/).
3. Another good link is [Estate Planning Resources and the Forest Legacy Program \(ct.gov\)](https://www.ct.gov/estateplanning/).

### Diversity

A healthy forest has a large diversity of native plant species, particularly trees that support a diverse array of fungi and wildlife (animals, insects, microbes). A healthy forest also features multiple layers of native vegetation, maximizing biodiversity and structural complexity. This means having trees of different ages, diameters, and heights. A healthy forest has standing dead trees (snags) and dead downed wood as essential habitat elements, which help retain moisture during droughts. A healthy forest is resilient because it can recover more effectively from diseases, pests, and extreme weather events. Increasing species and structural diversity of the forest provides multiple recovery pathways from disturbance.

To improve forest landscape diversity, consider establishing some patches of young forest. With over 89 birds, mammals, and reptiles that require young forest habitats, the young forest is nature's pantry for wildlife, thanks to its abundance of insects and berries. It also provides our mature woodland wildlife with important food and cover at critical times of the year. Ideally, 5-10% of our landscape should be young forest. Unfortunately, our landscape rarely has this much young forest because we prevent such forests from forming naturally via beaver activity, fires and floods. We can mimic natural disturbances with well-planned forest stewardship activities to create patches of young forest. Please visit [YoungForest.org](https://www.youngforest.org/) | [a fresh way to create wildlife habitat](#).

### Invasives/vines

Invasive species are typically from another part of the world, and when established here, they have no native predators or competitors to regulate their population. When left uncontrolled, they out-compete what would grow naturally, including tree regeneration and other native understory vegetation. Native understory growth supports a greater diversity of native insects and arthropods that wildlife needs to forage on. It also provides more nutrient-rich berries for birds. Exotic invasive understory growth can provide a better habitat for ticks and associated pathogens while significantly reducing biodiversity. Please see: [Home | Connecticut Invasive Plant Working Group \(uconn.edu\)](https://www.uconn.edu/invasive-plant-working-group/) and [https://www.conservect.org/wp-content/uploads/2021/06/Invasive\\_guide\\_2020\\_web.pdf](https://www.conservect.org/wp-content/uploads/2021/06/Invasive_guide_2020_web.pdf)

Control methods include mechanical and chemical processes. In a shady forest, cutting a vine is enough to kill it. Invasive shrubs are not so easy. Pulling the invasives out by the roots can be effective, but extremely difficult and labor-intensive. Yearly cutting back of the above-ground stems, during the growing season, will keep the invasives under control and may even kill them after a few years. The most effective control method is applying herbicide to the green foliage. You can also cut the invasive shrubs and treat the fresh stumps with herbicides to prevent resprouting.



Buckthorn Blaster herbicide applicator for vine and invasive shrub eradication: [Official Buckthorn Blaster® by NAISMA](https://www.naisma.com/)

### Lawns and fields

Fields offer opportunities to help pollinators and native insects. Insects, pollinators (including bees, butterflies,

moths, beetles, flies, wasps, and hummingbirds), and the many birds that depend on them, are in severe decline. By delaying annual mowing until after the first hard frost and before the start of spring plant growth, you will allow pollinators to use your fields as food and habitat during the growing season. Another habitat management strategy is to mow one-third to one-half each year on a rotational schedule, with a 6" or more cutting height. Insects will overwinter and nest in leaf litter, snags, dead wood and uncut plant stalks. This provides birds with much-needed food during the winter. For this reason, early spring mowing is best, before the green-up and bird-nesting season. Please remember that healthy meadows store more than double the carbon of a mowed lawn.

There are also many opportunities to create pollinator-friendly habitats/food sources by adding native plantings, allowing lawn areas to revert to natural areas, and leaving leaves and needles to cover the ground out to at least the drip line of the tree canopy. It is best to have a diversity of site-appropriate plants that bloom throughout the season (early, middle, and late), and to prioritize warm-season grasses and forbs as larval host plants rather than cool-season grasses (e.g., pasture grasses). Birds will pick through the winter leaves for insects. For more information, please visit: [Pollinator Pathway \(pollinator-pathway.org\)](https://www.audubon.org/pollinator-pathway). Link for native plantings: <https://www.audubon.org/native-plants>.

### **Boundaries**

Boundaries must be well-marked to protect the property from trespass and encroachment. Painted blazes are typically used to mark property boundaries. A blaze is a hand-sized shallow scrape in the bark. This scrape will last decades and not harm the tree if done correctly. When painted, this blaze is quite visible and long-lasting. Trees within arm's length of the boundaries are blazed, with the blazes facing the boundary line. Use only paint marks on the neighbor's side of the line, without blazes. The blazes should be repainted at least every ten years. Custom signs can also be hung at approximately 100-foot intervals using galvanized steel or aluminum nails. Keep the nails sticking out an inch from the tree so that the tree has room to grow without pushing the signs out. Understory vegetation and debris can be cleared along boundary lines to facilitate easy inspection. Please consider hiring a forester to locate and mark property boundaries.

### **Wildlife**

Your forest, and the State of Connecticut in general, is lucky to have a significant and diverse component of mature oak trees (mature trees have reached maximum height). Oak trees are considered a wildlife keystone species due to the large amount and diversity of life they support, surpassing that of any other tree. Acorns, especially white oak acorns, provide the most nutritious plant-based protein for almost 90 wildlife species. Oaks overwhelmingly host the most moth and butterfly caterpillars (over 500), which anchor a biodiverse food web. Oak forests have higher bird abundance and diversity than other forest types. Oaks produce the thickest, most ecologically beneficial, and longest-lasting leaf litter, accompanied by the most abundant and diverse soil biology. This top-of-the-line leaf litter can keep out invasive exotic stilt grass and jumping worms. It also purifies and holds the most water. For these reasons, preserving and encouraging oak growth and health in your forest is important.

Parts of this forest contain legacy trees, also known as old-field trees or wolf trees. These trees grew in open pastures as a source of shade for livestock before the current forest began to form. They are much older than the surrounding forest. Because they were once open-grown, they have large, spreading crowns and large branches low on the trunk. When the pastures were abandoned, they became a significant seed source for the present forest. These large, old trees are structurally complex, with numerous cavities, hollows, thick branches, and rough, thick bark. They are also prolific seed producers, including acorns and nuts. This structural complexity and prolific seed production attract a vast array of insects, birds, and mammals. Underground, the old trees are also the hub and source of the complex fungal soil mycorrhizal growth that all trees depend on for water and nutrients. To make them healthier and more vigorous, such legacy trees could be protected and perhaps even given more sunlight by cutting some surrounding trees. These agrarian vestiges have become the ecological hubs in your forest. They are also a great source of future large snags and ecologically valuable large dead, downed wood.

## Ecological Services and Climate Change Mitigation

Forests remove carbon dioxide from the atmosphere (a process called sequestration), produce oxygen, and remove many pollutants from the air and water. Forests absorb heavy rains and release that water to streams and underground aquifers during droughts. Forests regulate temperature and moisture levels. Forests provide homes for countless insects, including most of our many pollinators. In the spring, the tree canopy provides valuable bee pollen when the meadows, gardens and crops have yet to flower (particularly our oaks and maples). Forests store carbon in the below-ground roots and soil, as well as in above-ground vegetation, dead wood, and fallen leaves.

These services are enhanced by having a diverse mix of native tree species in various sizes, ages and arrangements. Forest diversity translates into habitat diversity, wildlife diversity, insect diversity and climate resilience. Sustainable, scientifically based forest management to remove forest products and promote young forests or regeneration of desired species has no long-term adverse effect on your forest's ability to provide these vital ecological services. When trees are young and growing fast, they sequester carbon at high rates, and once they are large (over 18" in diameter and often older), they store the most carbon. Whether you choose to manage your forest actively or not, your forest does an excellent service to our planet's health just by being a healthy forest.

Forests store carbon in different pools, and the amounts in these pools change over time. The pools comprise live aboveground elements (trees, shrubs, and other plants), live belowground elements (roots and fungi), deadwood (standing dead trees [snags] and downed logs), litter (leaves, needles, and small branches), and soil organic matter. Sequestration is the process by which forests remove carbon dioxide from the atmosphere, primarily via tree photosynthesis. A younger forest (10-60 years old) stores relatively little carbon but is likely at or near its peak sequestration rate. An older, more mature forest (60 years or older) stores more carbon, with a gradually slowing sequestration rate. A mix of sequestration and storage in multi-aged forests creates a resilient carbon profile. Please remember that using and harvesting local wood is an integral part of climate mitigation and a vital tool for improving the resilience of our forests to climate change.

## Mapping

Attached to this report is a geo-referenced map that the landowner can use with mapping apps. This digital map shows the landowner's location on the property, and the landowner can also record tracks and waypoints.

1. To get map layers and to view maps, please visit [CT ECO Home \(cteco.uconn.edu\)](http://cteco.uconn.edu).
2. To get soil maps and associated soil descriptions, please visit [Web Soil Survey - Home \(usda.gov\)](http://websoilsurvey.sc.egov.usda.gov) and follow the instructions on the first page of this website.
3. For instruction, please see [Tutorials | Center for Land Use Education and Research \(uconn.edu\)](http://www.ct.gov/landuse).

## Getting Assistance

Here is a list of every licensed forester in CT: <https://www.depdata.ct.gov/forestry/ForestPractitioner/directry.pdf>

Additionally, we are seeking applicants for the Master Woodland Manager certification (MWM). Here is a link for MWM: [Connecticut Master Woodland Manager Program | Connecticut Forest & Park Association \(ctwoodlands.org\)](http://ctwoodlands.org)

To learn more about the NRCS EQIP funding Program, click [NRCS EQIP Program Help](#)

Visit CT DEEP's grants website: [Grants and Financial Assistance](#)

Wildlife habitat projects: [Partners for Fish and Wildlife | U.S. Fish & Wildlife Service](#)

To prevent erosion and water quality issues: [Best Management Practices Manual](#)



---

APPENDIX OF LATIN NAMES

---

TREES

|                |                                |                  |                              |
|----------------|--------------------------------|------------------|------------------------------|
| Red Oak        | <i>Quercus rubra</i>           | Eastern Hemlock  | <i>Tsuga canadensis</i>      |
| Tulip-Poplar   | <i>Liriodendron tulipifera</i> | Aspen            | <i>Populus</i> sp.           |
| Red Maple      | <i>Acer rubrum</i>             | Blackgum         | <i>Nyssa sylvatica</i>       |
| Black Birch    | <i>Betula lenta</i>            | Yellow Birch     | <i>Betula alleghaniensis</i> |
| White Oak      | <i>Quercus alba</i>            | Chestnut Oak     | <i>Quercus montana</i>       |
| Hickory        | <i>Carya</i> Sp.               | Black Cherry     | <i>Prunus serotina</i>       |
| Black Oak      | <i>Quercus velutina</i>        | White Pine       | <i>Pinus strobus</i>         |
| Scarlet Oak    | <i>Quercus coccinea</i>        | Sassafras        | <i>Sassafras albidum</i>     |
| American Beech | <i>Fagus grandifolia</i>       | Paper Birch      | <i>Betula papyrifera</i>     |
| Sugar Maple    | <i>Acer saccharum</i>          | White Ash        | <i>Fraxinus americana</i>    |
| Sycamore       | <i>Platanus occidentalis</i>   | Eastern Redcedar | <i>Juniperus virginiana</i>  |
| Slippery Elm   | <i>Ulmus rubra</i>             | Basswood         | <i>Tilia americana</i>       |
| Pitch Pine     | <i>Pinus rigida</i>            | Gray Birch       | <i>Betula populifolia</i>    |

NATIVE UNDERSTORY

|                     |                                |                  |                             |
|---------------------|--------------------------------|------------------|-----------------------------|
| Eastern Hophornbeam | <i>Ostrya virginiana</i>       | Spicebush        | <i>Lindera benzoin</i>      |
| Musclewood          | <i>Carpinus caroliniana</i>    | Witch Hazel      | <i>Hamamelis virginiana</i> |
| Serviceberry        | <i>Amelanchier</i> Sp          | Mountain Laurel  | <i>Kalmia latifolia</i>     |
| Lowbush blueberry   | <i>Vaccinium angustifolium</i> | Huckleberry      | <i>Gaylussacia baccata</i>  |
| Highbush Blueberry  | <i>Vaccinium corymbosum</i>    | Sweet Pepperbush | <i>Clethra alnifolia</i>    |
| Striped Maple       | <i>Acer pensylvanicum</i>      | Holly            | <i>Smilax</i>               |
| Hobblebush          | <i>Viburnum lantanoides</i>    | Greenbrier       | <i>Ilex</i>                 |
| Maple-leaf Viburnum | <i>Viburnum acerifolium</i>    |                  |                             |

EXOTIC INVASIVES

|                  |                               |                |                              |
|------------------|-------------------------------|----------------|------------------------------|
| Barberry         | <i>Berberis</i> Sp.           | Burning Bush   | <i>Euonymus alatus</i>       |
| Multi-flora Rose | <i>Rosa multiflora</i>        | Bittersweet    | <i>Celastrus orbiculatus</i> |
| Privet           | <i>Ligustrum</i> Sp           | Honeysuckle    | <i>Lonicera</i> Sp           |
| Russian Olive    | <i>Elaeagnus angustifolia</i> | Tree-of-Heaven | <i>Ailanthus altissima</i>   |